


## Wagyu EBV Graph for ITOSHIGENAMI (IMP USA)

[Home](#)  
 [Animal Enquiry](#)  
 [EBV Enquiry](#)  
 [Mating Predictor](#)  
 [Member Enquiry](#)  
 [Sale Catalogues](#)  
 [Semen/Embryo Catalogues](#)  
 [Download Files](#)  
 [Online Transactions](#)  
 [Tag Orders](#)

EBV Percentiles for ITOSHIGENAMI (IMP USA)



[Switch Graph](#)  
 [Graph Explanation](#)

October 2021 Wagyu BREEDPLAN														
	Gestation Length (days)	Birth Wt (kg)	200 Day Wt (kg)	400 Day Wt (kg)	600 Day Wt (kg)	Mat Cow Wt (kg)	Milk (kg)	Scrotal Size (cm)	Carcase Wt (kg)	Eye Muscle Area (sq cm)	Rump Fat (mm)	Retail Beef Yield (%)	Marble Score	Marble Fineness (%)
EBV	+1.5	<b>-1.8</b>	-3	-7	-16	-9	-7	-1.7	-13	+3.4	+1.0	-0.2	<b>+2.3</b>	+0.33
<u>Accuracy</u>	98%	99%	99%	99%	99%	98%	99%	98%	98%	98%	98%	96%	98%	97%
Breed Avg. EBVs for 2019 Born Calves <a href="#">Click for Percentiles</a>														
EBV	+0.0	+1.0	+9	+15	+18	+21	+0	-0.2	+15	+1.4	-0.3	+0.1	+0.8	+0.16

**Traits Analysed:** Genomics

**Statistics:** Number of Herds: 130, Progeny Analysed: 3263, Scan Progeny: 726, Carcase Progeny: 473, Number of Dtrs: 1026

SELECTION INDEX VALUES		
Market Target	Index Value	Breed Average
Wagyu Breeder Index	+\$ 146	+\$ 142
Self Replacing Index	+\$ 188	+\$ 142
Fullblood Terminal Index	+\$ 159	+\$ 119
F1 Terminal Index	+\$ 173	+\$ 109

Estimated Breeding Values can only be directly compared to other EBVs calculated in the same analysis. Results from different analyses are likely based upon different datasets and different underlying parameters and trait definitions.

Information contained on this web database, including but not limited to pedigree, DNA information, Estimated Breeding Values (EBVs) and Index values, is based on data recorded on the society/association database which was supplied by members and/or third parties. Whilst every effort is made to ensure the accuracy of the information, the ABRI, the society/association, their officers and employees assume no responsibility for its content, use or interpretation. Data submitted to the database may have errors in it which can not be detected by current testing technology. For this reason, users ought to consider if they need to obtain independent testing of the relevant animal (if possible) to ensure the data is accurate.

BREEDPLAN results are calculated using software developed by the Animal Genetics and Breeding Unit, a joint venture of NSW Department of Primary Industries and the University of New England, which receives funding for this purpose from Meat and Livestock Australia Limited.